

An axially symmetric solution of metric-affine gravity

E.J. Vlachynsky*, R. Tresguerres†, Yu.N. Obukhov‡, and F.W. Hehl

Institute for Theoretical Physics, University of Cologne

D-50923 Köln, Germany

Abstract

We present an exact stationary *axially symmetric* vacuum solution of metric-affine gravity (MAG) which generalises the recently reported spherically symmetric solution. Besides the metric, it carries nonmetricity and torsion as post-Riemannian geometrical structures. The parameters of the solution are interpreted as mass and angular momentum and as dilation, shear and spin charges.

PACS no.: 04.50.+h; 04.20.Jb; 03.50.Kk

Typeset using REVTeX

*Permanent address: Department of Mathematics, University of Newcastle, Newcastle, NSW 2308, Australia

†Permanent address: Consejo Superior de Investigaciones Cientificas, Serrano 123, 28006 Madrid, Spain

‡Permanent address: Department of Theoretical Physics, Moscow State University, 117234 Moscow, Russia

The geometry of MAG (‘metric-affine gravity’) is described by the curvature two-form R_{α}^{β} , the nonmetricity one-form $Q_{\alpha\beta}$, and the torsion two-form T^{α} which are the gravitational field strengths for linear connection, metric and coframe, respectively. The corresponding physical sources are the canonical energy-momentum and hypermomentum three-forms. The latter includes the dilation, shear and spin currents associated to matter. The field equations and the formalism are comprehensively described in [1].

In the recent paper [2], we presented an exact spherically symmetric solution of MAG which has a mass and three nontrivial hypermomentum charges (dilation, shear and spin). Here we generalise this result and describe an axially symmetric solution.

We start from the Lagrangian [2],

$$\begin{aligned} V = \frac{1}{2\kappa} & \left[-a_0 R^{\alpha\beta} \wedge \eta_{\alpha\beta} - 2\lambda \eta + T^{\alpha} \wedge * \left(\sum_{I=1}^3 a_I {}^{(I)}T_{\alpha} \right) \right. \\ & + 2 \left(\sum_{I=2}^4 c_I {}^{(I)}Q_{\alpha\beta} \right) \wedge \vartheta^{\alpha} \wedge *T^{\beta} + Q_{\alpha\beta} \wedge * \left(\sum_{I=1}^4 b_I {}^{(I)}Q^{\alpha\beta} \right) \Big] \\ & - \frac{z_4}{2} R^{\alpha\beta} \wedge {}^{*(4)}Z_{\alpha\beta}, \end{aligned} \quad (1)$$

where $a_0, \dots, a_3, b_1, \dots, b_4, c_2, c_3, c_4, z_4$ are coupling constants. We choose the cosmological constant λ to be positive for an overall repulsion. As usual, the star $*$ denotes the Hodge dual, and κ is the Einstein gravitational constant. The metric signature is $(-+++)$. The fourth irreducible piece of the curvature reads ${}^{(4)}Z_{\alpha\beta} = R_{\gamma}^{\gamma} g_{\alpha\beta}/4$ (see [1] for details). Particular cases of (1) were considered earlier in [3–7].

Let us choose standard Boyer-Lindquist coordinates (t, r, θ, ϕ) . With the local Minkowski metric, $o_{\alpha\beta} = \text{diag}(-1, 1, 1, 1)$, we take as an ansatz the standard coframe for the Kerr solution with cosmological constant:

$$\begin{aligned} \vartheta^{\hat{0}} &= \sqrt{\frac{\Delta}{\Sigma}} \left(dt - j_0 \sin^2 \theta d\phi \right), \\ \vartheta^{\hat{1}} &= \sqrt{\frac{\Sigma}{\Delta}} dr, \\ \vartheta^{\hat{2}} &= \sqrt{\frac{\Sigma}{f}} d\theta, \\ \vartheta^{\hat{3}} &= \sqrt{\frac{f}{\Sigma}} \sin \theta \left[-j_0 dt + (r^2 + j_0^2) d\phi \right], \end{aligned} \quad (2)$$

where $\Delta = \Delta(r)$, $\Sigma = \Sigma(r, \theta)$, $f = f(\theta)$, and j_0 is a constant.

Similarly to [2], we assume that nonmetricity and torsion are represented by three one-forms (the Weyl covector, the third irreducible nonmetricity piece and the torsion trace) so that

$$Q_{\alpha\beta} = u(r, \theta) o_{\alpha\beta} \vartheta^{\hat{0}} + \frac{1}{9} v(r, \theta) \left(4\vartheta_{(\alpha} e_{\beta)} \rfloor \vartheta^{\hat{0}} - o_{\alpha\beta} \vartheta^{\hat{0}} \right), \quad (3)$$

$$T^\alpha = \frac{1}{3} \tau(r, \theta) \vartheta^\alpha \wedge \vartheta^{\hat{0}}. \quad (4)$$

Substituting the ansatz (2)-(4) into the field equations for the Lagrangian (1) (we refer the readers to [1] for the general form of the MAG field equations, (5.5.4)-(5.5.5), and to [2] for the detailed discussion of the model (1)) one finds:

$$u = k_0 N \frac{r}{\sqrt{\Delta \Sigma}}, \quad v = k_1 N \frac{r}{\sqrt{\Delta \Sigma}}, \quad \tau = k_2 N \frac{r}{\sqrt{\Delta \Sigma}}, \quad (5)$$

$$\Delta = r^2 + j_0^2 - 2\kappa M r - \frac{\lambda}{3a_0} r^2 (r^2 + j_0^2) + z_4 \frac{\kappa(k_0 N)^2}{2a_0}, \quad (6)$$

$$\Sigma = r^2 + j_0^2 \cos^2 \theta, \quad (7)$$

$$f = 1 + \frac{\lambda}{3a_0} j_0^2 \cos^2 \theta. \quad (8)$$

Here M and N are the integration constants which describe, respectively, the mass and the nonmetricity-torsion charges of the source. The coefficients k_0 , k_1 , k_2 are constructed in terms of the coupling constants,

$$k_0 = \left(\frac{a_2}{2} - a_0 \right) (8b_3 + a_0) - 3(c_3 + a_0)^2, \quad (9)$$

$$k_1 = -9 \left[a_0 \left(\frac{a_2}{2} - a_0 \right) + (c_3 + a_0)(c_4 + a_0) \right], \quad (10)$$

$$k_2 = \frac{3}{2} [3a_0(c_3 + a_0) + (8b_3 + a_0)(c_4 + a_0)], \quad (11)$$

and, as in [2], we find

$$b_4 = \frac{a_0 k + 2c_4 k_2}{8k_0}, \quad \text{with} \quad k := 3k_0 - k_1 + 2k_2. \quad (12)$$

The physical interpretation of the parameters of the solution as described above is clear: M and j_0 represent the mass and the angular momentum of Kerr type, while $(k_0 N)$, $(k_1 N)$,

and $(k_2 N)$ describe, respectively, the dilation (‘Weyl’) charge of the fourth irreducible non-metricity piece, the shear charge of the third irreducible nonmetricity piece and the spin charge of the second irreducible torsion part.

Work is in progress on the generalisation of our solution to the MAG version of the Plebański-Demiański family [8].

Acknowledgments This research was supported by the Deutsche Forschungsgemeinschaft (Bonn) under project He-528/17-1 and by the Graduate College “Scientific Computing” (Cologne-St.Augustin).

REFERENCES

- [1] F.W. Hehl, J.D. McCrea, E.W. Mielke, and Y. Ne'eman, Phys. Rep. **258** (1995) 1.
- [2] Yu.N. Obukhov, E.J. Vlachynsky, W. Esser, R. Tresguerres, and F.W. Hehl, *An exact solution of the metric-affine gauge theory with dilation, shear, and spin charges*, preprint of Inst. for Theor. Phys., University of Cologne (February 1996), to be submitted.
- [3] F.W. Hehl, E.A. Lord, and L.L. Smalley, Gen. Relat. Grav. **13** (1981) 1037.
- [4] V.N. Ponomariov and Yu. Obukhov, Gen. Relat. Grav. **14** (1982) 309.
- [5] R. Tresguerres, Z. Phys. **C65** (1995) 347.
- [6] R. Tresguerres, Phys. Lett. **A200** (1995) 405.
- [7] R.W. Tucker and C. Wang, Class. Quantum Grav. **12** (1995) 2587.
- [8] J.F. Plebański and M. Demiański, Ann. Phys. (NY) **98** (1976) 98.